

# The Role of Conservation Agriculture in Sustainable Agriculture

Peter R. Hobbs (PhD) Cornell University USA,  
Ken Sayre (PhD) CIMMYT, Mexico  
Raj Gupta (PhD) CIMMYT, India.

**Corresponding Author:** Peter R. Hobbs, Department of Crops and Soil Sciences, 609 Bradfield Hall, Cornell University, Ithaca NY 14850, USA. Phone: 607-254-7295; Fax: 607-255-2644; email: [ph14@cornell.edu](mailto:ph14@cornell.edu)

Requests for reprints same as corresponding author.

Word count without abstract or references: 10,681 words

Word count of Abstract: 295

**Keywords:** Cultivation, Conservation agriculture, Conservation tillage, Direct seeding, Zero-tillage, No-tillage, Rice-wheat systems, Bed planting, Mulching, Rotations.

## Abstract

The paper focuses on conservation agriculture (CA), defined as minimal soil disturbance (no-till) and permanent soil cover (mulch) combined with rotations, as a more sustainable cultivation system for the future. Cultivation and tillage, considered synonymous in this paper, play an important role in agriculture. The benefits of tillage in agriculture are explored before introducing conservation tillage (CT), a practice that was borne out of the American dust bowl of the 1930s. The paper then describes the benefits of CA, a suggested improvement on CT, where no-till, mulch and rotations significantly improve soil properties (physical, biological and chemical) and other biotic factors. The paper concludes that CA is a more sustainable and environmentally friendly management system for cultivating crops. Case studies from the rice-wheat areas of the Indo-Gangetic Plains of South Asia and the irrigated maize-wheat systems of NW Mexico are used to describe how CA practices have been used in these two environments to raise production sustainably and profitably. Benefits in terms of greenhouse gas emissions and their effect on global warming are also discussed. The paper concludes that agriculture in the next decade will have to sustainably produce more food from less land through more efficient use of natural resources and with minimal impact on the environment in order to meet growing population demands. This will be a tall order for agricultural scientists, extension personnel and farmers. Promoting and adopting CA management systems can help meet this goal.

# **The Role of Conservation Agriculture in Sustainable Agriculture**

## **Peter R. Hobbs, Ken Sayre and Raj Gupta**

### **Introduction**

Conservation agriculture (CA) defined (FAO CA web site) as minimal soil disturbance (no-till) and permanent soil cover (mulch) combined with rotations is a recent agricultural management system that is gaining popularity in many parts of the world. Cultivation is defined by the Oxford English dictionary as “the tilling of land”, “the raising of a crop by tillage” or “to loosen or break up soil”. Other terms used in this dictionary include “improvement or increase in (soil) fertility”. All these definitions indicate that cultivation is synonymous with tillage or ploughing.

The other important definition that has been debated and defined in many papers is the word “sustainable”. The Oxford dictionary defines this term as “capable of being borne or endured, upheld, defended, maintainable”. Something that is sustained is “kept up without intermission or flagging, maintained over a long period”. This is an important concept in today’s agriculture since the human race will not want to compromise the ability of its future offspring to produce their food needs by damaging the natural resources used to feed the population today.

This paper will introduce and promote CA as a modern agricultural practice that can enable farmers in many parts of the world to achieve the goal of sustainable agricultural production. But first the paper discusses some issues related to tillage.

### **Cultivation techniques or tillage**

The history of tillage dates back many millennia when humans changed from hunting and gathering to more sedentary and settled agriculture mostly in the Tigris, Euphrates, Nile, Yangste and Indus river valleys (Hillel, 1991). Reference to ploughing or tillage is found from 3000 BC in Mesopotamia (Hillel, 1998). Lal, 2001 explains the historical development of agriculture with tillage being a major component of management practices. With the advent of the industrial revolution in the 19<sup>th</sup> century, mechanical power and tractors became available to undertake tillage operations; today an array of equipment is available for tillage and agricultural production. The following summarizes the reasons for using tillage:

1. Tillage was used to soften the soil and prepare a seedbed that allowed seed to be placed easily at a suitable depth into soil moisture using seed drills or manual equipment. This results in good, uniform seed germination.
2. Wherever crops grow, weeds also grow and compete for light, water and nutrients. Every gram of resource used by the weed is one less gram for the crop. By tilling their fields farmers were able to shift the advantage from the weed to the crop and allow the crop to grow without competition early in its growth cycle with resulting higher yield.
3. Tillage helped release soil nutrients needed for crop growth through mineralization and oxidation after exposure of soil organic matter to air.
4. Previous crop residues were incorporated along with any soil amendments (fertilizers, organic or inorganic) into the soil. Crop residues, especially loose residues, create problems for seeding equipment by raking and clogging.

5. Many soil amendments and their nutrients are more available to roots if they incorporated into the soil; some nitrogenous fertilizers are also lost to the atmosphere if not incorporated.
6. Tillage gave temporary relief from compaction by using implements that could shatter below ground compaction layers formed in the soil.
7. Tillage was determined to be a critical management practice for controlling soil borne diseases and some insects.

There is no doubt that this list of tillage benefits was beneficial to farmers, but at a cost to him and the environment and the natural resource base on which farming depended. The utility of ploughing was first questioned by a forward looking agronomist in the 1930s, Edward H. Faulkner, in a manuscript called "Ploughman's Folly" (Faulkner, E.H., 1943). In a foreword to a book entitled "Ploughman's folly and a second look" by EB Faulkner, Paul Sears notes that:

*"Faulkner's genius was to question the very basis of agriculture itself -- the plough. He began to see that the curved moldboard of the modern plough, rather than allowing organic matter to be worked into the soil by worms and other burrowing animals, instead buries this valuable material under the subsoil where it remains like a wad of undigested food from a heavy meal in the human stomach"*(Faulkner, E.B. 1987)

The tragic dust storms in the mid-western United States in the 1930s was a wake up call to how man's interventions in soil management and ploughing led to un-sustainable agricultural systems. In the 1930s it was estimated that 91 Mha of land was degraded by severe soil erosion (Utz *et al.*, 1938); this area has been dramatically reduced today.

### **Conservation tillage and conservation agriculture**

Since the 1930s and during the next 75 years members of the farming community have been advocating a move to reduced tillage systems that use less fossil fuel, reduce runoff and erosion of soils and reverse the loss of soil organic matter. The first 50 years was the start of the conservation tillage movement and today a large percentage of agricultural land is cropped using these principles. However, in the book "No-Tillage Seeding" (Baker *et al.*, 2002; a new edition of this excellent book will be available in late 2005) explains "As soon as the modern concept of reduced tillage was recognized, everyone, it seems, invented a new name to describe the process." The book goes on to list 14 different names for reduced tillage along with rationales for using these names. The book is also an excellent review of the mechanization and equipment needs of no-tillage technologies. Baker (2002) defines conservation tillage as:

*"Conservation tillage is the collective umbrella term commonly given to no-tillage, direct-drilling, minimum-tillage and/or ridge-tillage, to denote that the specific practice has a conservation goal of some nature. Usually, the retention of 30% surface cover by residues characterizes the lower limit of classification for conservation-tillage, but other conservation objectives for the practice include conservation of time, fuel, earthworms, soil water, soil structure and nutrients. Thus residue levels alone do not adequately describe all conservation tillage practices"* (Baker *et al.*, 2002)

This has led to confusion among the agricultural scientists and more important the farming community. To add to the confusion, the term “*conservation agriculture*” has recently been introduced by FAO (Food and Agriculture Organization web site) and others and its goals defined by FAO as follows:

*“Conservation agriculture (CA) aims to conserve, improve and make more efficient use of natural resources through integrated management of available soil, water and biological resources combined with external inputs. It contributes to environmental conservation as well as to enhanced and sustained agricultural production. It can also be referred to as resource efficient or resource effective agriculture”* (FAO)

This obviously encompasses the “*sustainable agricultural production*” need that all mankind obviously wishes to achieve. But this term is often not distinguished from conservation tillage. FAO mentions in its CA website that:

*“Conservation tillage is a set of practices that leave crop residues on the surface which increases water infiltration and reduces erosion. It is a practice used in conventional agriculture to reduce the effects of tillage on soil erosion. However, it still depends on tillage as the structure forming element in the soil. Never the less, conservation tillage practices such as zero tillage practices can be transition steps towards Conservation Agriculture.”*

In other words conservation tillage uses some of the principles of conservation agriculture but has more soil disturbance.

### **Conservation agriculture defined**

FAO has characterized conservation agriculture as follows:

*“Conservation Agriculture maintains a permanent or semi-permanent organic soil cover. This can be a growing crop or dead mulch. Its function is to protect the soil physically from sun, rain and wind and to feed soil biota. The soil micro-organisms and soil fauna take over the tillage function and soil nutrient balancing. Mechanical tillage disturbs this process. Therefore, zero or minimum tillage and direct seeding are important elements of CA. A varied crop rotation is also important to avoid disease and pest problems.”* (FAO web site)

Data reported by Derpsch, 2005 indicates that the extent of no-tillage adoption worldwide is just over 95 million hectares. This figure is used as a proxy for CA although not all of this land is permanently no-tilled or has permanent ground cover. Table 1 details the extent of no-tillage by country worldwide. Six countries have more than one million hectares. South America has the highest adoption rates and has more permanent no-till and permanent soil cover. Both Argentina and Brazil had significant lag periods to reach 1 million hectares in the early 1990s and then expanded rapidly to the present day figures of 18.3 and 23.6 mha, respectively for these countries. By adopting the no-tillage system Derpsch (2005) estimates that Brazil

increased its grain production by 67.2 million tons in 15 years with additional revenue of 10 billion dollars. Derpsch also estimates that at an average rate of 0.51 t ha<sup>-1</sup>yr<sup>-1</sup> Brazil sequestered 12 million tons of carbon on 23.6 million hectares of no-tillage land. Tractor use is also significantly reduced saving millions of litres of diesel.

**Table 1: Extent of no-tillage adoption worldwide**

<b>Country</b>	<b>Area under No-tillage (mha) 2004/2005</b>
USA	25.30
Brazil	23.60
Argentina	18.27
Canada	12.52
Australia	9.00
Paraguay	1.70
Indo-Gangetic-Plains (**)	1.90
Bolivia	0.55
South Africa	0.30
Spain	0.30
Venezuela	0.30
Uruguay	0.26
France	0.15
Chile	0.12
Colombia	0.10
China	0.10
Others (Estimate)	1.00
<b>Total</b>	<b>95.48</b>

Source: Derpsch, 2005.

\*\* Includes area in India, Pakistan, Bangladesh and Nepal I South Asia

The three key principles of conservation agriculture are permanent residue soil cover, minimal soil disturbance and crop rotations. FAO recently added controlled traffic to this list. Each of these will be briefly dealt with before providing some case studies. Table 2 shows a comparison of CA with CT and traditional tillage (TT).

### **Permanent or semi-permanent organic soil cover**

Unger *et al.*, 1988 reviews the role of surface residues on water conservation and indicates that this association between surface residues, enhanced water infiltration and evaporation led to the adoption of CT after the 1930s dust bowl problem. Research since that time has documented beyond doubt the importance of surface residues on soil water conservation and reduction in wind and water erosion (Unger *et al.*, 1988; Van Doran & Allmaras, 1978). Allmaras *et al.*, 1991 reviews much of the literature on conservation tillage up to that time and goes on to describe a whole array of conservation tillage-planting systems operational in the US; their adoption and benefits.

Table 2. A comparison of tillage, conservation tillage (CT) and conservation agriculture (CA) for various issues.

<b>Issues</b>	<b>Traditional tillage TT</b>	<b>Conservation tillage CT</b>	<b>Conservation agriculture CA</b>
Practice	Disturbs the soil and leaves a bare surface	Reduces the soil disturbance in TT and keeps the soil covered	Minimal soil disturbance and soil surface permanently covered
Erosion	Wind and soil erosion maximum	Wind and soil erosion reduced significantly	Wind and soil erosion the least of the three
Soil physical health	The lowest of the 3	Significantly improved	The best practice of the 3
Compaction	Used to reduce compaction but can also induce it by destroying biological pores.	Reduced tillage is used to reduce compaction	Compaction can be a problem but use of mulch and promotion of biological tillage helps reduce this problem.
Soil biological health	The lowest of the 3 because of frequent disturbance	Moderately better soil biological health	More diverse and healthy biological properties and populations
Water infiltration	Lowest after soil pores clogged	Good water infiltration	Best water infiltration
Soil organic matter	Oxidises soil organic matter and causes its loss	Soil organic buildup possible in the surface layers	Soil organic buildup in the surface layers even better than CT.
Weeds	Controls weeds but also causes more weed seed to germinate	Reduced tillage controls weeds but also exposes other weed seed for germination	Weeds are a problem especially in the early stages of adoption, but problems are reduced with time and residues can help suppress weed growth
Soil temperature	Surface soil temperature more variable	Surface soil temperature intermediate in variability	Surface soil temperature moderated the most.
Diesel use and costs	Diesel use high	Diesel use intermediate	Diesel use much reduced
Costs production	Highest costs	Intermediate costs	Lowest costs
Timeliness	Operations can be delayed	Intermediate timeliness of operations	Timeliness of operations more optimal
Yield	Can be lower where planting delayed	Yields same as TT	Yields same as TT but can be higher if planting done more timely

This paper will not go into detail about other soil conserving practices that are utilized throughout the world and over time, like terracing or contour bunds that essentially are designed to prevent soil erosion on sloping lands. Lal, 2001 describes some of these systems and notes that the effectiveness of these systems depends on proper construction and regular maintenance; if not done properly degradation can be catastrophic.

Kumar & Goh, 2000 review the effect of crop residues and management practices on soil quality, soil nitrogen dynamics and recovery and crop yield. The review concludes that crop residues of cultivated crops are a significant factor for crop production through their affects on soil physical, chemical and biological functions as well as water and soil quality. They can

have both positive and negative effects and the role of agricultural scientists is to enhance the positive effects.

This paper will restrict the discussion of crop residues to their benefits when used as mulch. Crop residues result when a previous crop is left anchored or loose after harvest or when a cover crop (legume or non-legume) is grown and killed or cut to provide mulch. Externally applied mulch in the form of composts and manures can also be applied, although the economics of transport of this bulky material to the field may restrict its use to higher value crops like vegetables.

The energy of raindrops falling on a bare soil result in destruction of soil aggregates, clogging of soil pores and rapid reduction in water infiltration with resulting runoff and soil erosion.. Mulch intercepts this energy and protects the surface soil from soil aggregate destruction, enhances the infiltration of water, and reduces the loss of soil by erosion. (Freebairn & Boughton, 1985; McGregor *et al.*, 1990; Dormaar & Carefoot, 1996). Topsoil losses of 46.5 t/ha have been recorded with conventional tillage on sloping land after heavy rain in Paraguay compared to 0.1 t/ha under no-till cultivation (Derpsch and Moriya, 1999). No-till plus mulch reduces surface soil crusting, increases water infiltration, reduces runoff and gives higher yield than tilled soils (Cassel *et al.*, 1995; Thierfelder *et al.*, 2005). Similarly, the surface residue, anchored or loose, protects the soil from wind erosion (Michels *et al.*, 1995). The dust bowl is a useful reminder of the impacts of wind and water erosion when soils are left bare.

Surface mulch helps reduce water losses from the soil by evaporation and also helps moderate soil temperature. This promotes biological activity and enhances nitrogen mineralization, especially in the surface layers (Dao, 1993; Hatfield & Prueger, 1996). This is a very important factor in tropical and sub-tropical environments but has been shown to be a hindrance in temperate climates due to delays in soil warming in the spring and delayed germination (Schneider & Gupta, 1985; Kaspar *et al.*, 1990; Burgess *et al.*, 1996; Swanson & Wilhelm, 1996). Fabrizzi *et al.*, 2005 showed that no-till had lower soil temperatures in the Spring in Argentina, but conventional tillage had higher maximum temperatures in the summer and that average temperatures during the season were similar.

Karlen *et al.*, 1994 showed that normal rates of residue combined with zero-tillage resulted in better soil surface aggregation and that this could be increased by adding more residues. Recent papers confirm this observation; Madari *et al.*, 2005 show no-till (NT) with residue cover had higher aggregate stability, higher aggregate size values and total organic carbon in soil aggregates than conventional tillage (CT) in Brazil; Roldan *et al.*, 2003 after 5 years NT had increased soil enzymes, soil organic carbon (SOC), microbial biomass (MBM) and wet aggregate stability compared CT; soil quality improved with more biomass. They conclude that NT is a sustainable technology.

A cover crop and the resulting mulch or previous crop residue help reduce weed infestation through competition and not allowing weed seeds the light often needed for germination. There is also evidence of allelopathic properties of cereal residues in respect to inhibiting surface weed seed germination (Lodhi & Malik, 1987; Steinsiek *et al.*, 1982; Jung *et al.*, 2004). Weeds will be controlled when the cover crop is cut, rolled flat or killed. Farming

practice that maintains soil micro-organisms and microbial activity can also lead to weed suppression by the biological agents (Kennedy 1999).

Cover crops contribute to the accumulation of organic matter in the surface soil horizon (Roldan *et al.*, 2003; Alvear *et al.*, 2005; Riley *et al.*, 2005; Madari *et al.*, 2005; Diekow *et al.*, 2005) and this effect is increased when combined with NT. Mulch also helps with recycling of nutrients, especially when legume cover crops are used, through the association with below-ground biological agents and by providing food for microbial populations. Greater carbon and nitrogen were reported under no-tillage and conservation tillage compared to ploughing (Campbell *et al.*, 1995, 1996a and b). Others have shown that this is restricted to the surface horizons and that the reverse occurs at greater depths in humid soils of eastern Canada (Anger *et al.*, 1997). Schultz, 1992, showed C and N declined by 6% with burning but increased by 1% with stubble retention. Vagen *et al.*, 2005 conclude that the largest potential for increasing soil organic carbon is through the establishment of natural or improved fallow systems (agroforestry) with attainable C accumulation rates from 0.1 to 5.3 Mg C ha<sup>-1</sup> yr<sup>-1</sup>. They continue to say that in cropland, addition of crop residues or manure in combination with no-till can yield attainable C accumulation rates up to 0.36 Mg C ha<sup>-1</sup> yr<sup>-1</sup>. Soil organic carbon (SOC) is a key indicator of soil quality and Lal, 2005 calculates that increasing SOC by 1 Mg ha<sup>-1</sup> yr<sup>-1</sup> can increase food grain production by 32 million Mg yr<sup>-1</sup> in developing countries. Heenan *et al.*, 2004 in Australia show that changes in SOC at the surface ranged from a loss 8.2 t ha<sup>-1</sup> for continuous tilled cereals and residues burnt to a gain of 3.8 t ha<sup>-1</sup> where stubble was retained and soil no-tilled. Nitrogen content followed similar trends. If the rotation included a legume, SOC accumulation was the highest.

Soil microbial biomass (SMB) has commonly been used to assess below ground microbial activity and is a sink and source for plant nutrients. Amendments such as residues and manures promote, while burning and removal of residues decrease SMB (Collins *et al.*, 1992; Heenan *et al.*, 2004; Doran 1980; Angers *et al.*, 1993a and b; Alvear *et al.*, 2005). Balota *et al.*, 2004 in Brazil in a 20 year experiment showed that residue retention and NT increased total C by 45% and SMB by 83% at 0-50 cm depth compared to CT. Soon & Arshad, 2005 showed SMB was greater with NT than CT by 7-36%; frequent tillage resulted in a decrease in both total and active MBC. Increased SMB occur rapidly in a few years following conversion to reduced tillage (Ananyeva *et al.*, 1999; Alvarez & Alvarez, 2000). Increased microbial biomass increased soil aggregate formation, increased nutrient cycling through slow release of organically stored nutrients and also assisted in pathogen control (Carpenter-Boggs *et al.*, 2003).

Cover crops help promote biological soil tillage through their rooting ; the surface mulch provides food, nutrients and energy for earthworm, arthropods and micro-organisms below-ground that also biologically till soils. Use of deep-rooted cover crops and biological agents (earthworms, etc.) can also help to relieve compaction under zero-tillage systems. There is a lot of literature that looks at burning, incorporation and removal of crop residues on soil properties and much less where mulch is left on the surface. An early paper by McCalla, 1958 showed that bacteria, actinomycetes, fungi, earthworms and nematodes were higher in residue mulched fields than those where the residues were incorporated. More recent studies also show more soil fauna in no-tillage, residue retained management treatments compared to tillage plots

(Buckerfield & Webster, 1996; Nuutinen, 1992; Karlen *et al.*, 1994; Hartley *et al.*, 1994; Riley *et al.*, 2005; Birkas *et al.*, 2004; Kemper & Derpsch, 1981; Clapperton, 2003). Tillage disrupts and impairs soil pore networks including those of mycorrhizal hyphae, an important component for phosphorus availability in some soils (Evans & Miller, 1990; McGonigle & Miller 1996). Zero-tillage thus results in a better balance of microbes and other organisms and a healthier soil.

Groundcover promotes an increase in biological diversity below but also above-ground; the number of beneficial insects was higher where there was groundcover and mulch (Jaipal *et al.*, 2002; Kendall *et al.*, 1995) and these help keep insect pests in check.

Interactions between root systems and rhizobacteria effect crop health, yield and soil quality. Release of exudates by plants activate and sustain specific rhizobacterial communities that enhance nutrient cycling, nitrogen fixing, bio-control of plant pathogens, plant disease resistance and plant growth stimulation. Sturz & Christie, 2003 give a review of this topic. Groundcover would be expected to increase biological diversity and increase these beneficial effects.

### **Minimal Soil Disturbance**

Many of the benefits of minimal soil disturbance were mentioned in the above section on permanent soil cover and in fact combining these two practices is important for obtaining the best results. The following comparisons between tillage and zero-tillage systems are made to highlight some other benefits not mentioned above.

Tractors consume large quantities of fossil fuels that add to costs while also emitting greenhouse gases (mostly CO<sub>2</sub>) and contributing to global warming when used for ploughing (Grace *et al.*, 2003). Animal based tillage systems are also expensive since farmers have to maintain and feed a pair of animals for a year for this purpose. Animals also emit methane, a greenhouse gas 21 times more potent for global warming than carbon dioxide (Grace *et al.*, 2003). Zero-tillage reduces these costs and emissions. Farmer surveys in Pakistan and India show that zero-till of wheat after rice reduces costs of production by US\$60 per hectare mostly due to less fuel (60-80 liters ha<sup>-1</sup>) and labor (Hobbs & Gupta, 2004).

Tillage takes valuable time that could be used for other useful farming activities or employment. Zero-tillage minimizes time for establishing a crop. The time required for tillage can also delay timely planting of crops, with subsequent reductions in yield potential (Hobbs & Gupta 2003). By reducing turnaround time to a minimum, zero-tillage can get crops planted on time and thus increase yields without greater input cost. Turnaround time in this rice-wheat system from rice to wheat varies from 2 to 45 days since 2-12 passes of a plough is used by farmers to get a good seedbed (Hobbs & Gupta, 2003). With zero-till wheat this time is reduced to just one day.

Tillage and current agricultural practices result in decline of soil organic matter due to increased oxidation over time, leading to soil degradation, loss of soil biological fertility and resilience (Lal, 1994). Although this SOM mineralization liberates nitrogen and can lead to improved yields over the short term, there is always some mineralization of nutrients and loss

by leaching into deeper soil layers. This is particularly significant in the tropics where organic matter reduction is processed more quickly, with low soil carbon levels resulting after only one or two decades of intensive soil tillage. Zero-tillage, on the other hand, combined with permanent soil cover, has been shown to result in a build-up of organic carbon in the surface layers (Campbell *et al.*, 1996a; Lal, 2005). No-tillage minimizes SOM losses and is a promising strategy to maintain or even increase soil C and N stocks (Bayer *et al.*, 2000).

Although tillage does afford some relief from compaction, it is itself a major cause of compaction, especially when repeated passes of a tractor are made to prepare the seedbed or to maintain a clean fallow. Zero-tillage reduces dramatically the number of passes over the land and thus compaction. However, natural compaction mechanisms and the one pass of a tractor mounted zero-till drill will also result in compaction. The FAO CA web site now includes “controlling in-field traffic” as a component of conservation agriculture; this is accomplished by having field-traffic follow permanent tracks. This can also be accomplished by using a ridge-till or permanent bed planting system rather than planting on the flat (Sayre & Hobbs, 2004). Some farmers feel that sub-soiling or chiseling may be needed to resolve below ground compaction layers before embarking on a no-till strategy, especially in dryer areas.

Higher bulk densities and penetration resistance have been reported under zero-tillage compared to tillage (Gantzer & Blake, 1978), and is described as natural for zero-tillage (Ehlers & Claupein 1994). This problem is more in soils with low stability soil aggregates (Ehlers *et al.*, 1983). Bautista *et al.*, 1996 working in a semi-arid ecosystem found that zero-tillage plus mulch reduced bulk density. The use of zero-till using a permanent residue cover even when bulk density (BD) was higher, resulted in higher infiltration of water in no-till systems (Sayre and Hobbs, 2004; Shaver *et al.*, 2002). Scientists hypothesized that continued use of reduced, shallow and zero-tillage would require a shift to short term, conventional tillage to correct soil problems. However, Logsdon & Karlen, 2004 show that BD is not a useful indicator and confirm that farmers need not worry about increased compaction when changing from CT to NT on deep loess soils in USA. Fabrizzi *et al.*, 2005 working in Argentina had higher BD and penetration resistance in NT but the values were below thresholds that could affect crop growth; wheat yields were the same as tilled plots. This experiment left residues on the surface in NT. The authors concluded that the experiment had a short time frame and more time was needed to assess the effect of BD.

The role of tillage on soil diseases is discussed by Leake (2003) with examples of the various diseases affected by tillage. He concludes that the role of tillage on diseases is unclear and acknowledges that a healthy soil with high microbial diversity does play a role by being antagonistic to soil pathogens. He also suggests that no-till farmers need to adjust management to control diseases through sowing date, rotation, and resistant cultivars to help shift the advantage from the disease to the crop. A list of the impacts of minimum tillage on specific crops and their associated pathogens can be found in Sturz *et al.*, 1997.

An added, economic consideration is that tillage results in more wear-and-tear on machinery and higher maintenance costs for tractors than under zero-tillage systems.

## **Rotations**

Crop rotation is an agricultural management tool with ancient origins. Howard (1996) reviewed the cultural control of plant diseases from an historical view and includes examples of disease control through rotation. The rotation of different crops with different rooting patterns combined with minimal soil disturbance in zero-till systems promotes a more extensive network of root channels and macro-pores in the soil. This helps in water infiltration to deeper depths. Because rotations increase microbial diversity, the risk of pests and disease outbreaks from pathogenic organisms is reduced, since the biological diversity helps keep pathogenic organisms in check (Leake, 2003). The discussion of the benefits of rotations will be handled in other chapters of this publication.

Integrated pest management (IPM) should also be added to the conservation agriculture set of recommendations, since if one of the requirements is to promote soil biological activity, minimal use of toxic pesticides and use of alternative pest control methods that do not affect these critical soil organisms is needed. A review of IPM in CA can be found in Leake, 2003.

## **Equipment for Conservation Agriculture**

Before going on to describe a couple of case studies from Asia and Mexico, there is a need to discuss the critical importance of equipment for success with CA; zero-till and conservation agriculture is bound to fail if suitable equipment is not available to drill seed into residues at the proper depth for good germination. It is urgent that CA equipment is perfected, available and adopted for this new farming system. Iqbal *et al.*, 2005 studied NT under dryland conditions in Pakistan and showed NT gave lower yields than CT, but the experiment was planted with improper equipment and with no mention of residue management; the results are therefore suspect.

There are some excellent reviews of the equipment needs for zero-tillage systems. Baker *et al.*, 2002 devotes an entire book to this topic and will soon publish a new edition to this useful publication in 2005. A book on Conservation Agriculture (Environment, Farmer Experiences, Innovations and Socio-economic Policy) edited by Garcia-Torres *et al.*, 2003 has several papers on equipment for small scale and large-scale farmers. The main requirements of equipment in a conservation agriculture system is a way to handle loose straw (cutting or moving aside), seed and fertilizer placement, furrow closing and seed/soil compaction. There is also a need for small scale farmers to adapt direct drill seeding equipment to animal or small tractor power sources (reduced weight and draft requirements) and reduce costs so equipment is affordable by farmers, although use of rental and service providers allow small scale farmers to use this system even if they do not own a tractor or seeder.

A simple 3 row small grain seeder has been developed for small scale animal powered farmers in Bolivia (Wall *et al.*, 2003). This equipment uses a shovel rather than a disc opener to save weight. It has straw wheels attached to the coulter to help move residues aside and reduce clogging. It also has the benefit that it can be used in ploughed or unploughed soil. The main benefit farmers mentioned about this drill was savings in time; it takes 10 hours to plant a hectare with this machinery and 12 days for the conventional tillage and seeding method. The conventional system also required the farmer to walk 100 km ha<sup>-1</sup> to undertake all the tillage and seeding with his animals. The stand with the new drill was 246 +/- 37 plants/ m<sup>2</sup> compared

to 166 +/- 39 for the conventional system. The cost of the drill was only \$330-390 in Bolivia. Similar information is provided by Ribeiro (2003) for planters in Brazil. These can be manually applied jabber planters to animal drawn planters. In both countries the participation of farmers, local manufacturers and extensionists were vital for success.

Saxton & Morrison, 2003 look at equipment needs for large-scale farmers where tractor horsepower, weight and draft are less important. Earlier machines were developed for clean tilled farm fields, whereas new machines provide precision seed placement through consistent soil penetration and depth and also supply fertilizer in bands which is crucial for minimizing nutrient losses in zero-till systems. This paper discusses the use of disc openers versus hoe and chisel openers and the use of additional straw and chaff spreading devices.

### **Case study from the Rice-Wheat Systems of South Asia**

The first case study looks at the 13.5 million hectares of the rice-wheat systems of the Indo-Gangetic Plains for South Asia (RWC web site). The traditional cultivation technique used for growing rice in this system, but also in much of the rice growing regions of Asia, is done by wet ploughing of soils in the main rice field (puddling), followed by transplanting of rice seedlings grown in separate seed beds. Interestingly, this system of rice cultivation is often cited as being used for centuries without declining productivity. However, that was at relatively low, subsistence rice yield levels. There are several excellent reviews of a number of Asian long term experiments using modern varieties on this issue, some rice-wheat and others rice-rice, with some showing yield declines while others do not (Dawe *et al.*, 2000; Cassman *et al.*, 1995; Abrol *et al.*, 1997).

Puddling was done by farmers over the centuries for very specific reasons, the most important being to help control weeds; farmers found that keeping soils anaerobic and flooded reduced weed problems; but also hand weeding was easier with these softened soils. The puddling essentially reduced water percolation and infiltration and ponded the water on the surface. Less is written on the puddling effect on soil biological properties although some work is available from research done at IRRI in the 1990s (Reichardt *et al.*, 1997; Reichardt *et al.*, 2001). The authors conclude that soil microbial biomass plays a significant role as a passive nutrient pool and suggests that its reduction found in puddled soils in the second half of the cropping season could be a mechanism that contributes to declining productivity in continuous rice cropping systems. Little has been published on soil microbes in rice-wheat systems.

When modern varieties of wheat and rice were introduced to South Asia in the 1960s, farmers in NW India and Pakistan introduced rice into their wheat systems and farmers in the Eastern side of S. Asia introduced wheat into their rice systems; wheat was grown in the cool dry season and rice in the warm, wet monsoon months. This intensified the system that has grown to 13.5 million hectares since the 1960s. It is now one of the most important cropping systems for food security in S. Asia along with rice-rice systems. One of the main issues that confronted farmers when this new system was introduced and found feasible and profitable, was the soil physical properties left after harvest of a puddled, transplanted rice crop. The effect of puddling reduced soil structure, especially stable soil aggregates, and led to formation of compaction layers (Hobbs *et al.*, 1993). Soil cracking was higher under intensive puddling (Mohanty *et al.*, 2004). Unpuddled direct seeded rice maintained the soil

in a better physical condition, although yields were lower where weeds were not controlled. Farmers ploughed their fields many times to obtain a suitable seed bed for planting wheat (Hobbs & Gupta, 2003, table 7.1). This ploughing takes time and often results in late planting and decline in wheat yield potential plus many other negative effects (Hobbs & Gupta, 2003; Hobbs & Gupta, 2004).

The solution for late planting and problems of delayed turnaround from rice harvest to wheat planting came from the introduction of zero-tilled wheat into rice stubbles that started in the region in the mid-1980s. Efforts to adapt and promote resource conserving technologies (RCT's that include NT) in the Indo-Gangetic Plains (IGP) have been underway for nearly 3 decades but it is only in the past 4 to 5 years that the technologies are finding accelerated acceptance by the farmers (figure 1). The spread of NT is taking place in the irrigated RW regions but are yet to be rooted in the rainfed agro-ecoregions. In the last 2004-05 wheat season, it was estimated that nearly 2 million hectares of zero-till wheat was being grown by 425,000 farmers in the 4 South Asian countries (RWC Highlights 2005); both big and small scale farmers adopted this technology with small scale farmers renting zero-till drill services from service providers. The key to this rapid adoption in the last 5 years was the use of farmer participatory approaches to allow farmers to experiment with the technology in their own fields and promotion of the local machinery manufacturers in the region to be partners in the program; cheap, affordable, effective drills are available based on the use of the inverted-T coulter technology that was introduced into India and Pakistan from New Zealand.

One major need of this system is the development and availability of equipment that will allow good germination of rice and wheat while at the same time minimizing soil disturbance and sowing the seed and banded fertilizer into loose and anchored stubbles. The RWC members are working vigorously in partnership with local manufacturers and farmers to make new equipment available for experimentation at an affordable price, with provisions for after sales service and supply of needed spare parts to make this system successful. Recently, multi-crop zero-till ferti-seed drills fitted with inverted -T openers, disk planters, punch planters, trash movers or roto-disk openers have been developed for seeding into loose residues (RWC Highlights 2004-05) (figure2).

Many advantages have been mentioned and characterized for this innovation including US\$145 million savings in fuel costs (2004 costs) and the benefits of less greenhouse gas emissions, less weeds, more beneficial insect activity, improved water use efficiency, but also important, higher yields at less cost, improved incomes from wheat and savings in time that can be used for other productive uses (Hobbs & Gupta, 2004). It is anticipated that as machinery manufacturers keep pace with demand for drills, the message about the benefits of zero-till wheat reaches more farmers and farmer mindsets about the need for tillage are overcome that the acreage for this innovation will cover the bulk of the wheat planted after rice in South Asia.

However, the story does not end here. In order for the benefits of zero-tilled wheat to be seen in the entire system and for soil physical and biological health to improve and the rice-wheat system to become more sustainable, rice practices will also have to change. A move is already afoot in the rice-wheat consortium partners (national and international programs) to

research, experiment and promote a move to zero-till direct seeded rice; a more aerobic rice system that does away with puddling of soils. These aerobic systems are based on direct seeded rice systems either on the flat or raised beds, with and without tillage and with or without transplanting (RWC DSR paper). The ultimate system would allow farmers to grow zero-till rice followed with zero-till wheat. Wheat yields are best after direct seeded rice without puddling (DSNP) (Tripathi *et al.*, 2005; Hobbs *et al.*, 2002).

Various national and international research and breeding agencies are now exploring aerobic rice (Bouman *et al.*, 2002). The main issue to resolve relates to effective weed control in a non-puddled rice system. Various innovative, integrated ways are being sought to handle this problem including use of cover crops and mulches, more competitive rice varieties against weeds and use of selective herbicides. Availability of roundup ready transgenic rice and/or development of cultivars suited to direct seeding with zero-till drills, having early vigor, and competitive with weeds would go a long way to help resolve this issue. One encouraging technology intercropped direct seeded rice with a green manure (*Sesbania aculeata*). After a month, the crop is sprayed with 2,4-D herbicide to knock down the green manure and kill any germinating broadleaf weeds. The dieing weeds and GM provide nutrients to the rice crops as they decompose but new weeds are suppressed by the ground cover and allelopathic properties of the mulch. The results look good and this research will be reported soon. In addition to zero-till rice and wheat, attempts are also being made to diversify the cropping systems by introducing other crops into new rotations that help break disease and insect cycles and provide more income and diversity for farmers. This may help with some other problems that have surfaced when rice is shifted from anaerobic to aerobic systems. Widespread infestations of the root knot nematode (*Meloidyogyne graminicola*) on rice were found when direct seeded rice was grown instead of puddled rice in Bangladesh (Padgham *et al.*, 2004).

### **Case Study from Mexico**

Maize-wheat cropping patterns are common in the irrigated NW areas of Mexico and the rainfed areas of the altiplano areas of central Mexico. In both situations, the major limiting factor is moisture. This case study introduces the concept of permanent bed systems for addressing efficient use of water. In bed systems, soil is raised in a ridge and furrow configuration. These bed systems involved tillage to prepare the soil before making the beds. However, many traditional bed planting systems did not receive tillage; the “*chinampas*” of pre-Colombian Mexico and the *waru warus* of Peru and Bolivia used crop residue mulching or only superficial tillage (Thurston, 1992). Bed systems reduce compaction in the rooting zone by confining wheel traffic to the furrow bottoms. The case study described here from Mexico looks at the results of using a permanent bed planting maize-wheat system where soil disturbance is minimized, crop residues are retained on the surface from previous crops and reshaping of beds is done only as needed between crop cycles (Limon-Ortega *et al.* 2002).

The experiment used for this paper was undertaken by the International Maize and Wheat Improvement Center (CIMMYT) in the State of Sonora in NW Mexico. Farmers have 225,000 hectares of irrigated land in this area with maize, sorghum, soybean, safflower, dry beans, cotton, and wheat the major crops. Ninety five percent of the farmers now grow crops on beds with farmers changing from the conventional planting on the flat with basin irrigation in the past 20 years (Aquino 1998). This change was a result of water shortages from the water

storage reservoir system; farmers had to find more efficient water use systems in order to expand acreage. Results suggest that bed planted systems need 29% less water than flat planting systems for an 8% higher yield (Sayre and Hobbs, 2004). Most of the farmers still use conventional tillage to remake the beds for each new crop, but results that are reported below suggest that use of permanent bed systems where tillage is minimized and crop residues are left on the surface will be more sustainable.

The treatments in this long term trial were as follows:

1. Conventional tillage with formation of new beds for each crop and with all the crop residues incorporated.
2. Permanent beds with all the residues burned
3. Permanent beds with 30% of the residues retained on the surface and the rest baled for fodder
4. Permanent beds with the maize residue baled for fodder and the wheat residue retained
5. Permanent beds with all the residues retained.

A detailed account of this experiment can be found in Sayre and Hobbs, 2004. Yield differences were small in the first 5 years supporting the idea that some transitional years are needed before changes occur in soil properties with changes in management. But differences started to appear in the 6<sup>th</sup> year with the permanent bed treatment with all residues retained the best treatment; the worst plot, with significantly less yield was the permanent bed with residues burnt. The treatment with conventional tillage with residues incorporated was statistically at par with the best permanent bed system but incurred higher costs for land preparation.

Organic matter, nitrogen levels, surface soil aggregate size, and soil microbial biomass were higher in the permanent beds with residue retention. One valuable insight from this experiment was the lower soil strength/compaction in all treatments, except the one where residues were burned. The addition of the residue plays a significant role in reducing compaction at the soil surface and increasing water infiltration in minimal tilled plots. Similar data confirmed this finding in a similar maize-wheat long term experiment under rainfed conditions in the altiplano areas near Mexico City (Govaerts et al., 2005). In this rainfed experiment, zero-tilled plots with residue retention resulted in higher and more stable yields than conventionally tilled plots with residues incorporated. Zero-tilled plots without residue retention had much reduced yields. In the same experiment, permanent raised beds combined with rotation and residue retention yielded the same as zero-tilled plots with residue retention. The bed system gave farmers the added advantage of being able to use more varied weed and fertilizer practices.

Larger scale demonstrations have been planted on farmer fields. The permanent beds averaged  $7.2 \text{ t ha}^{-1}$  compared to  $6.2 \text{ t ha}^{-1}$  for the conventionally made beds. The data also show that average returns over variable costs increased by 75% for the permanent bed system with residue retention compared to the conventional tilled treatment. The importance of suitable equipment that will allow seeding of crops into permanent beds with residue retention cannot be over emphasized. Systems based on discs, punch planters and strip tillage are being experimented with in Mexico and in South Asia (Sayre and Hobbs, 2004).

### **Climate change and conservation agriculture**

Lal, 2005 suggests that by adopting improved management practices on agricultural land (use of no-till and crop residues) food security would be enhanced but it would also offset fossil fuel emissions at the rate of  $0.5 \text{ Pg C yr}^{-1}$ . Climate change is likely to strongly affect rice-wheat, rice-rice and maize-based cropping systems that today account more for than 80 % of the total cereals grown on more than 100 million ha of agricultural lands in South Asia. Global warming may be beneficial in some regions, but harmful in those regions where optimal temperatures already exist; an example would be the rice-wheat mega-environments in the Indo-Gangetic plains that account for 15% of global wheat production. Agronomic and crop management practices have to aim at reducing  $\text{CO}_2$  and other greenhouse gas emissions by reducing tillage and residue burning and improving nitrogen use efficiency. In the Indo-Gangetic plains, resource-conserving technologies continue to expand in the rice-wheat cropping systems and save 50-60 L of diesel  $\text{ha}^{-1}$  plus labor, and significantly reduce release of  $\text{CO}_2$  to the environment. Methane emissions that have a warming potential 21 times that of  $\text{CO}_2$  are common and significant in puddled, anaerobic paddy fields and also when residues are burnt. This GHG emission can be mitigated by shifting to an aerobic, direct seeded or NT rice system. A review of the other benefits of direct seeding and NT in RW areas of South Asia can be found in Grace *et al.*, 2003. Nitrous oxide has 310 times the warming potential of carbon dioxide and its emissions are affected by poor nitrogen management. Sensor based technologies for measuring normalized differential vegetative index (NVDI) and moisture index have been used in Mexico and S. Asia to help improve the efficiency of applied nitrogen and reduce nitrous oxide emissions.

### **Conclusions**

Crop production in the next decade will have to produce more food from less land by making more efficient use of natural resources and with minimal impact on the environment. Only by doing this will food production keep pace with demand and the productivity of land preserved for future generations. This will be a tall order for agricultural scientists, extension personnel and farmers. Use of productive but more sustainable management practices described in this paper can help resolve this problem. Crop and soil management systems that help improve soil health parameters (physical, biological and chemical) and reduce farmer costs are essential. Development of appropriate equipment to allow these systems to be successfully adopted by farmers is a pre-requisite for success. Overcoming traditional mindsets about tillage by promoting farmer experimentation with this technology in a participatory way will help accelerate adoption. Encouraging donors to support this long term, applied research with sustainable funding is also an urgent requirement.

### **References**

Abrol I.P, Bronson K.F, Duxbury J.M, & Gupta R.K. (eds.). 1997. *Long-term soil fertility experiments in rice-wheat cropping systems*: Proceedings of a Workshop, 15-18 Oct 1996, Surajkund, Haryana, India. New Delhi, India, Consortium Paper Series 1. Rice-Wheat Consortium for the Indo-Gangetic Plains, New Delhi, India. pp 23.

Allmaras RR, Langdale GW, Unger PW, Dowdy RH, & Van Doren DM. 1991. Adoption of conservation tillage and associated planting systems. In: Lal R, Pierce FJ, editors. *Soil management for sustainability*. Papers from a workshop in Edmonton, Alberta, August 1989. Soil and Water Conservation Society. USA. p. 53-83.

Alvarez CR & Alvarez R. 2000. Short term effects of tillage systems on active soil microbial biomass. *Biol. Fert. Soils* **31**: 157-161.

Alvear M, Rosas A, Rouanet JL & Borie F. 2005. Effects of three soil tillage systems on some biological activities in an Ultisol from Southern Chile. *Soil & Tillage Research*. **82**: 195-202.

Ananyeva ND, Demkina TS, Jones WJ, Cabrera ML & Steen WC. 1999. Microbial biomass in soils of Russia under long term management practices. *Biol. Fert. Soils* **29**: 291-299.

Angers DA, Bissonette N, & Legere A. 1993a. Microbial and biochemical changes induced by rotation and tillage in a soil under barley production. *Can. J. Soil. Sci.* **73**: 39-50.

Angers DA, N'dayegamiye A, & Cote D. 1993b. Tillage-induced differences in organic matter of particulate size fractions and microbial biomass. *Soil Science Society of America Journal* **57**:512-516.

Angers DA, Bolinder MA, Carter MR, Gregorich EG, Drury CF, Liang BC, Voroney RP, Simard RR, Donald RG, Beyaert RP, & Martel J. 1997. Impact of tillage practices on organic carbon and nitrogen storage in cool, humid soils of eastern Canada. *Soil Tillage Res.* **41**: 191-201.

Aquino P. 1998. *The adoption of bed planting of wheat in the Yaqui Valley, Sonora, Mexico*. CIMMYT Wheat Special Report No. 17a. CIMMYT, Mexico, DF.

Baker CJ, Saxton KE, & Ritchie WR. 2002. *No-tillage seeding: science and practice*. 2<sup>nd</sup> Edition. CAB International. Oxford, UK.

Balota EL, Filho AC, Andrade DS & Dick RP. 2004. Long-term tillage and crop rotation effects on microbial biomass and C and N mineralization in a Brazilian Oxisol. *Soil & Tillage Research* **77**: 137-145.

Bautista S, Bellot J & Ramon-Vallejo V. 1996. Mulching treatment for post fire soil conservation in semi-arid eco-systems. *Arid Soil Res. Rehabilitation* **10**: 235- 242.

Bayer C, Mielniczuk J, Amado TJC, Martin-Neto L & Fernandes SV. 2000. Organic matter storage in a sandy loam Acrisol affected by tillage and cropping systems in southern Brazil. *Soil & Tillage Research* **54**: 101-109.

Birkas M, Jolankai M, Gyuricza C & Percze A. 2004. Tillage effects on compaction, earthworms and other soil quality indicators in Hungary. *Soil & Tillage Research* **78**: 185-196.

Bissett MJ & O'Leary GJ. 1996. Effects of conservation tillage on water infiltration in two soils in south-eastern Australia. *Australian Journal of Soil Research* **34**: 299-308.

Bouman BAM, Hengsdijk H, Hardy B, Bindraban PS, Tuong TP & Ladha JK. (Eds.) 2002. *Water-wise rice production*. Proceedings of the International Workshop on Water-wise rice production, 8-11 April 2002, Los Banos, Philippines. Los Banos (Philippines): International Rice Research Institute. 356 p.

Buckerfield JC, & Webster KA. 1996. Earthworms, mulching, soil moisture and grape yields: Earthworm responses to soil management practices in vineyards, Barossa Valley, South Australia. *Aust. N.Z. Wine Ind. J.* **11**: 47-53.

Burgess MM, Mehuys GR, & Madramootoo CA. 1996. Tillage and crop residue effects on corn production in Quebec. *Agron. J.* **88**: 792-797.

Campbell CA, McConkey BG, Zentner RP, Dyck FB, Selles F, & Curtin D. 1995. Carbon sequestration in a Brown Chernozem as affected by tillage and rotation. *Can. J. Soil Sci.* **75**: 449-458.

Campbell CA, McConkey BG, Zentner RP, Selles F, & Curtin D. 1996a. Long-term effects of tillage and crop rotations on soil organic C and N in a clay soil in South-Western Saskatchewan. *Can. J. Soil Sci.* **76**: 395-401.

Campbell CA, McConkey BG, Zentner RP, Dyck FB, Selles F, & Curtin D. 1996b. Tillage and crop rotation effects on soil organic C and N in a coarse-textured Typic Haploboroll in South-Western Saskatchewan. *Soil Tillage Res.* **37**: 3-14.

Carpenter-Boggs L, Stahl PD, Lindstrom MJ, Schumacher TE. 2003. Soil microbial properties under permanent grass, conventional tillage, and no-till management in South Dakota. *Soil & Tillage Research.* **71**: 15-23.

Cassel DK, Raczkowski CW & Denton HP. 1995. Tillage effects on corn production and soil physical conditions. *Soil Science Society of America Journal* **59**:1436-1443.

Cassman KG, De Datta SK, Olk DC, Alcantera J, Samson M, Descalsota JP, & Dixon M. 1995. Yield decline and the nitrogen economy of long-term experiments on continuous, irrigated rice systems in the tropics. In: Lal R, Stewart BA, (Eds). *Soil Management: Experimental Basis for Sustainability and Environmental Quality*. C&R Press, Boca Raton, Florida. p. 181-222.

Clapperton MJ. 2003. Increasing soil biodiversity through conservation agriculture: Managing the soil as a habitat. *Proceedings of the Second World Congress on Conservation Agriculture: Producing in Harmony with Nature*, Iguassu Falls, Parana-Brazil, August 11-15, 2003. Food and Agriculture Organization, Rome [CD].

Collins HP, Rasmussen PE, & Douglas CL Jr. 1992. Crop rotation and residue management effects on soil carbon and microbial dynamics. *Soil Science Society of America Journal* **56**: 783-788.

Dao TH. 1993. Tillage and winter wheat residue management effects on water infiltration and storage. *Soil Science Society of America Journal*. **57**: 1586-1595.

Dawe D, Dobermann, A, Moya P, Abdulrachman S, Bijay Singh, Lal, P, Li SY, Lin B, Panaullah G, Sariam, O, Singh, Y, Tan PS, & Zhen QX. 2000. How widespread are yield declines in long-term rice experiments. *Field Crops Research* **66**: 175-193.

Derpsch R, & Moriya K. 1999. Implications of soil preparation as compared to no-tillage on the sustainability of crop production: Experiences from South America. In: Reddy MV, (Ed.) *Management of Tropical Agro-Ecosystems and the Beneficial Soil Biota*. Science Publishers, Enfield, NH, p. 49-65.

Derpsch R. 2005 The extent of Conservation Agriculture adoption worldwide: Implications and impact. *Proceedings of the Third World Congress on Conservation Agriculture: Linking Production, Livelihoods and Conservation.*, Nairobi, Kenya, , October 3-7, 2005. [CD].

Diekow J, Mielniczuk J, Knicker H, Bayer C, Dick DP & Kogel-Knabner I. 2005. Soil C and N stocks as affected by cropping systems and nitrogen fertilization in a southern Brazil Acrisol managed under no-tillage for 17 years. *Soil & Tillage Research* **81**: 87-95.

Dormaar JF, & Carefoot JM. 1996. Implication of crop residue and conservation tillage on soil organic matter. *Can. J. Plant Sci.* **76**: 627-634.

Ehlers W, Kopke U, Hess F, & Bohm W. 1983. Penetration resistance and root growth of soils in tilled and untilled loess soil. *Soil & Tillage Research*. **3**: 261-275.

Evans DG & Miller MH. 1990. The role of external mycelial network in the effect of soil disturbance upon vesicular-arbuscular micorrhizal colonization of young maize. *New Phytol* **114**: 65-71.

Fabrizzi KP, Garcia FO, Costa JL & Picone LI. 2005. Soil water dynamics, physical properties and corn and wheat responses to minimum and no-tillage systems in the southern Pampas of Argentina. *Soil & Tillage Research* **81**: 57-69.

FAO conservation agriculture web site:

<http://www.fao.org/waicent/faoinfo/agricult/ags/AGSE/agsee/general/OBJECT.htm>

Faulkner EH. 1943. *Plowman's folly*. Michael Joseph Ltd. London, England.

Faulkner EB. 1987. *Plowman's folly and a second look*. Island Press. Washington DC.

Freebairn DM & Boughton WC. 1985. Hydrologic effects of crop residue management practices. *Aust. J. Soil Res.* **23**: 23-55. 1985.

Gantzer CJ & Blake GR. 1978. Physical characteristics of a La Seur clay loam following no-till and conventional tillage. *Agronomy Journal* **70**: 853-857.

Garcia-Torres L, Benites J, Martinez-Vilela A, & Holgado-Cabrera A. (editors). 2003. *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-economy, Policy*. Kluwer Academia Publishers. Dordrecht/Boston/London.

Govaerts B, Sayre KD & Deckers J. 2005. Stable high yields and zero tillage and permanent bed planting? *Field Crops Research*. **94 (1)**: 33-42.

Grace PR, Harrington L, Jain MC & Robertson GP. 2003. Long-term sustainability of the tropical and subtropical rice-wheat system: An environmental perspective. In J.K. Ladha, J. Hill, R.K. Gupta, J. Duxbury and R.J. Buresh. (eds). *Improving the productivity and sustainability of rice-wheat systems: issues and impact*. ASA, Spec. Publ. 65, chapter 7: 27-43. ASA Madison, WI. USA

Hartley MJ, Ragman A, & Popay AJ. 1994. Use of mulches and herbicide in an apple orchard. *Proc. N.Z. Plant Prot. Conf.* **47**: 320-324.

Hatfield KL, & Pruegar JH. 1996. Microclimate effects of crop residues on biological processes. *Theor. Appl. Climatol.* **54**: 47-59.

Heenan DP, Chan KY & Knight PG. 2004. Long-term impact of rotation, tillage and stubble management on the loss of soil organic carbon and nitrogen from a Chromic Luvisol. *Soil & Tillage Research* **76**: 59-68.

Hillel. 1991. *Out of the Earth: Civilization and the life of the soil*. Free Press. New York.

Hillel. 1998. *Environmental soil physics*. Academic Press. San Diego.

Hobbs, P. R., T. Woodhead, & C. Meisner. 1993. Soil Physical factors important in the productivity of the rice-wheat rotation. In: Saunders DA, and Hettel G. (Eds.). *Wheat in the Heat Stressed Environments: Irrigated, Dry areas, and Rice-Wheat Farming Systems*. Proceedings of a workshop held in Nashipur, Bangladesh. February 14-15, 1993 CIMMYT Mexico,D.F.

Hobbs, P.R., Y. Singh, G.S. Giri, J.G. Lauren, and J.M. Duxbury. 2002. Direct seeding and reduced tillage options in the rice-wheat systems of the Indo-Gangetic Plains of South Asia. In S. Pandey, M. Mortimer, L. Wade, T.P Tuong, K. Lopez and B. Hardy. *Direct seeding: Research strategies and Opportunities. Proceedings of the international workshop on direct seeding in Asian rice systems: strategic research issues and opportunities*. 25-28 January, 2000, Bangkok. Thailand. International Rice Research Institute, Los Banos, Philippines. pp. 201-215.

Hobbs, P.R. & R.K. Gupta. 2003. Resource Conserving Technologies for Wheat in Rice-Wheat Systems. In J.K. Ladha, J. Hill, R.K. Gupta, J. Duxbury and R.J. Buresh. (eds) *Improving the productivity and sustainability of rice-wheat systems: issues and impact*. ASA, Spec. Publ. 65, paper 7: 149-171. ASA Madison, WI. USA

Hobbs, P.R. & R.K. Gupta. 2004. Problems and Challenges of No-Till Farming for the Rice-Wheat Systems of the Indo-Gangetic Plains in South Asia. In R. Lal, P. Hobbs, N. Uphoff & D.O. Hansen. (eds). *Sustainable Agriculture and the Rice-Wheat System*. Paper 6:101-119. Ohio State University. Columbus, Ohio, USA. Marcel Dekker, Inc., 270 Madison Avenue, New York, New York 10016 p. 101-121.

Howard RJ. 1996. Cultural control of plant diseases: a historical perspective. *Canadian Journal Plant Pathology* 18: 145-150.

Iqbal MM, Akhter J, Mohammad W, Shah SM, Nawaz H & Mahmood K. 2005. Effect of tillage and fertilizer levels on wheat yield, nitrogen uptake and their correlation with carbon isotope discrimination under rainfed conditions in north-west Pakistan. *Soil & Tillage Research* 80: 47-57

Jaipal S, Singh S, Yadav A, Malik RK, & Hobbs PR. 2002. Species diversity and population density of macro-fauna of rice-wheat cropping habitat in semi-arid subtropical northwest India in relation to modified tillage practices of wheat sowing. In: Malik RK, Balyan RS, Yadav A & Pahwa, SK. (Eds). 2002. *Herbicide-Resistance Management and Zero-Tillage in the Rice-Wheat Cropping System*: Proceedings of an International Workshop, Hissar, India, 4–6 March 2002. CCS Haryana University, Hissar, India. p. 166-171.

Jung WS, Kim KH, Ahn JK, Hahn SJ & Chung IM. 2004. Allelopathic potential of rice (*Oryza sativa* L.) residues against *Echinochloa crus-galli*. *Crop Protection* 23: 211-218.

Karlen DL, Wollenhaupt NC, Erbach DC, Berry EC, Swan JB, Eash NS, & Jordahl JL. 1994. Crop residue effects on soil quality following 10-years of no-till corn. *Soil Tillage Research*. 31: 149-167.

Kaspar TC, Erbach DC & Cruse RM. 1990. Corn response to seed-row residue removal. *Soil Sci. Soc. Am. J.* 54: 1112-1117.

Kemper B & Derpsch R. 1981. Results of studies made in 1978 and 1979 to control erosion by cover crops and no-tillage techniques in Paraná, Brazil. *Soil & Tillage Research* 1: 253-267.

Kendall DA, Chin NE, Glen DM, Wiltshire CW, Winstone L & Tidboald C. 1995. Effects of soil management on cereal pests and their natural enemies. In: Glen DM, Greaves MP & Anderson HM (Eds). *Ecology and Integrated Farming Systems* : Proceedings of the 13th Long Ashton International Symposium Chichester [England] ; New York : Wiley Press.

Kennedy AC. 1999. Soil microorganisms for weed management. *J. Crop Prod.* 2: 123-138.

Kumar K & Goh KM. 2000. Crop residues and management practices: effects on soil quality, soil nitrogen dynamics, crop yield and nitrogen recovery. *Advances in Agronomy* **68**: 198-279.

Lal R. 1994. Sustainable land use systems and resilience. In: D.J. Greenland and I. Szabolcs (eds). *Soil resilience and sustainable land use: proceedings of a symposium held in Budapest, 28 September to 2 October 1992, including the Second Workshop on the Ecological Foundations of Sustainable Agriculture (WEFSA II)*. CAB International. Wallingford, Oxon, U.K.

Lal R. 2001. Managing world soils for food security and environmental quality. *Advances in Agronomy*. **74**: 155-192.

Lal R. 2005. Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. *Land Degradation & Development*. (in press).

Leake AR. 2003. Integrated pest management for conservation agriculture. In: Garcia-Torres L, Benites J, Martinez-Vilela A, Holgado-Cabrera A. (editors). *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-economy, Policy*. Kluwer Academia Publishers. Dordrecht/Boston/London. p. 271-279.

Limon-Ortega A, Sayre KD, Drijber RA, & Francis CA. 2002. Soil attributes in a furrow-irrigated bed planting system in northwest Mexico. *Soil & Tillage Research* **63**: 123-132.

Lodhi MAK, & Malik KA. 1987. Allelopathy in agroecosystems: wheat phytotoxicity and its possible role in crop rotation. *Journal of Chemical Ecology* 13:1881-1891.

Logsdon S & Karlen DL. 2004. Bulk density as a soil indicator during conversion to no-tillage. *Soil & Tillage Research*. **78**: 143-149.

Madari B, Machado PLOA, Torres E, de Andrade AG & Valencia LIO. 2005. No tillage and crop rotation effects on soil aggregation and organic carbon in a Rhodic Ferralsol from southern Brazil. *Soil & Tillage Research* **80**: 185-200.

McCalla TM. 1958. Microbial and related studies of stubble mulching. *J. Soil Water Conserv.* **13**: 255-258

McGonigle TP & Miller MH 1996. Mycorrhizae, phosphorus adsorption and yield of maize in response to tillage. *Soil Sci. Soc. Am J.* 60: 1856-1861.

McGregor KC, Bengtson RL, & Mutchler CK. 1990. Surface and incorporated wheat straw effects on interrill runoff and soil erosion. *Transactions of the American Society of Agricultural Engineers*. **33**: 469-474.

Michels K, Sivakumar MVK, & Allison BE. 1995. Wind erosion control using crop residue. I. Effects on soil flux and soil properties. *Field Crops Research*. **40**: 101-110.

Mohanty M, Painuli DK & Mandal KG. 2004. Effect of puddling intensity on temporal variation in soil physical conditions and yield of rice (*Oryza sativa*) in a vertisol of central India. *Soil and Tillage Research* **76**: 83-94.

Nuutinen V. 1992. Earthworm community responses to tillage and residue management on different soil types in southern Finland. *Soil & Tillage Research* **23**: 221-239.

Padgham JL, Abawi GS, Duxbury JM, & Mazid MA. 2004. Impact of wheat on *Meloidogyne graminicola* populations in the rice-wheat system of Bangladesh. *Nematropica* **34**: 183-190.

Reichardt W, Mascarina G, Padre B & Doll J. 1997. Microbial communities of continuously cropped irrigated rice fields. *Applied and environmental microbiology*. **63** (1): 233-238.

Reichardt W, Brioes A, de Jesus R, & Padre B. 2001. Microbial population shifts in experimental rice systems. *Applied soil ecology*: **17**: 151-163.

Ribeiro RF. 2003. No-tillage equipment for small farms in Brazil. In: Garcia-Torres L, Benites J, Martinez-Vilela A, & Holgado-Cabrera A. (editors). *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-economy, Policy*. Kluwer Academic Publishers. Dordrecht/Boston/London. p. 263-271.

Rice-Wheat Consortium web site: <http://www.rwc.cgiar.org/index.asp>

RWC Highlights 2004-05: [http://www.rwc.cgiar.org/Pub\\_Info.asp?ID=113](http://www.rwc.cgiar.org/Pub_Info.asp?ID=113)

RWC Direct seeded rice (DSR) paper: [http://www.rwc.cgiar.org/Pub\\_Info.asp?ID=123](http://www.rwc.cgiar.org/Pub_Info.asp?ID=123)

Riley HCF, Bleken MA, Abrahamsen S, Bergjord AK & Bakken AK. 2005. Effects of alternative tillage systems on soil quality and yield of spring cereals on silty clay loam and sandy loam soils in cool, wet climate of central Norway. *Soil & Tillage Research* **80**: 79-93.

Roldan A, Caravaca F, Hernandez MT, Garcia C, Sanchez-Brito C, Velasquez M, Tiscareno M. 2003. No-tillage, crop residue additions, legume cover cropping effects on soil quality characteristics under maize in Patzcuaro watershed (Mexico). *Soil & Tillage Research* **72**: 65-73.

Saxton KE & Morrison JE. 2003. Farm machines for large-scale conservation agriculture. In: Garcia-Torres L, Benites J, Martinez-Vilela A, & Holgado-Cabrera A. (editors). *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-economy, Policy*. Kluwer Academia Publishers. Dordrecht/Boston/London. p. 255-263.

Sayre KD & Hobbs PR. 2004. The Raised-Bed System of Cultivation for Irrigated Production Conditions. *In* R. Lal, P. Hobbs, N. Uphoff & D.O. Hansen. (eds). *Sustainable*

*Agriculture and the Rice-Wheat System*. Ohio State University. Columbus, Ohio, USA. Paper 20: 337-355.

Schneider EC & Gupta SC. 1985. Corn emergence as influenced by soil temperature, matric potential and aggregate size distribution. *Soil Sci. Soc. Am. J.* **49**: 415-422.

Shaver TM, Peterson GA, Ahuja LR, Westfall DG, Sherrod LA, & Dunn G. 2002. Surface soil properties after twelve years of dryland no-till management. *Soil Science Society of America Journal.* **66**: 1292-1303. 2002.

Shultz JE. 1992. The effect of rotations, stubble treatments and nitrogen fertilizer on soil properties and crop production. *Natl. Soil Conf.*, 4<sup>th</sup> Abstract. 1992. p. 82.

Soon YK & Arshad MA. 2005. Tillage and liming effects on crop and labile soil nitrogen in an acid soil. *Soil & Tillage Research* **80**: 23-33.

Steinsiek JW, Oliver LR, & Collins F. 1982. Allelopathic potential of wheat (*Triticum aestivum*) straw on selected weed species. *Weed Science* 30:495-497.

Sturz AV, Carter MR & Johnston HW. 1997. A review of plant disease, pathogen interactions and microbial antagonism under conservation tillage in temperate humid agriculture. *Soil & Tillage Research.* **41**: 169-189.

Sturz AV & Christie BR. 2003. Beneficial microbial allelopathies in the root zone: the management of soil quality and plant disease with rhizobacteria. *Soil & Tillage Research* **72**: 107-123.

Swanson SP & Wilhelm WW. 1996. Planting date and residue rate effects on growth, partitioning and yield of corn. *Agron. J.* **88**: 205-210.

Thierfelder C, Amezquita E, & Stahr K. 2005. Effects of intensifying organic manuring and tillage practices on penetration resistance and infiltration rate. *Soil & Tillage Research* **82**: 211-226.

Thurston HD. 1992. *Sustainable practices for plant disease management in traditional farming systems*. Westview press. Boulder Colorado, USA.

Tripathi RP, Sharma P & Singh S. 2005. Tilth index: an approach to optimize tillage in rice-wheat system. *Soil & Tillage Research* **90**: 125-137.

Unger PW, Langdale DW, & Papendick RI. 1988. Role of crop residues – improving water conservation and use. In: Hargrove WL (Ed). *Cropping strategies for efficient use of water and nitrogen*. Special publication 51. American Society of Agronomy, Madison, Wisconsin. p. 69-100.

Utz EJ, Kellogg CE, Reed EH, Stallings JH & Munns EN. 1938. The problem: The nation as a whole. In: *Soils and Men. Yearbook of Agriculture*. USDA. Washington. DC. p. 84-110.

Vagen TG, Lal R, & Singh BR. 2005. Soil carbon sequestration in Sub-Saharan Africa: A review. *Land Degradation & Development*. **16**: 53-71.

Van Doran DM Jr. & Allmaras, RR. 1978. Effect of residue management practices on soil physical environment, microclimate and plant growth. In: Oschwald WR (Ed). *Crop residue management systems*. ASA special publication 31. American Society of Agronomy, Madison Wisconsin. p. 49-83.

Wall PC, Zambrana L, Gamez P, Sims B, & Calissaya A. 2003. Development of an animal drawn zero-tillage seeder for small grains. In: Garcia-Torres L, Benites J, Martinez-Vilela A, & Holgado-Cabrera A. (Eds). *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-economy, Policy*. Kluwer Academia Publishers. Dordrecht/Boston/London. p. 243-249.




Figure 1. Estimated area growth of no-till wheat in the Indo-Gangetic Plains for the past 10 years



Figure 2: Various equipment for planting wheat no-till in RWC. a) inverted-T coultter; b) Indian no-till drill using inverted T; c) disk type planter; d) star-wheel punch planter e) “Happy planter” where straw picked up and blown behind seeder; f) disk planter with trash mover.